





BOVINE RHINOTRACHEITIS — VIRUS DIARRHEA PARAINFLUENZA3 — RESPIRATORY SYNCYTIAL VIRUS VACCINE, MODIFIED LIVE VIRUS, MANNHEIMIA HAEMOLYTICA PASTEURELLA MULTOCIDA BACTERIN-TOXOID



PREVENTING THE PREVENTABLE WITH AN EFFECTIVE COMBINATION.

An effective combination vaccine that helps protect cattle against some of the most common—and costly—viral and bacterial pathogens associated with Bovine Respiratory Disease (BRD).



INDICATIONS:

Shown to be effective for the vaccination of healthy cattle five months of age or older against Infectious Bovine Rhinotracheitis, Bovine Viral Diarrhea Type 1a, Bovine Viral Diarrhea Type 2, Parainfluenza₃ and Bovine Respiratory Syncytial Virus, and *Mannheimia haemolytica* and *Pasteurella multocida*.

INSTRUCTIONS:

Aseptically rehydrate the freeze-dried modified live vaccine with the accompanying bottle of bacterin-toxoid. Mix reconstituted vial well. Administer 2.0 mL subcutaneously in the side of the neck followed by a second dose of *Mannheimia haemolytica-Pasteurella multocida* (PMH BACTM) to be given 14 to 28 days after the first dose.

CAUTIONS:

Do not vaccinate pregnant cows. Do not use in calves nursing pregnant cows. Abortions may result. Store at 2° to 8° C. Use entire contents when first opened. Inactivate unused contents before disposal. Do not vaccinate within 21 days of slaughter.

