## **Bovine Rhinotracheitis-**Parainfluenza<sub>3</sub> Vaccine Modified Live Virus






**Cattle Vaccine** 

50 Dose / 100 mL

## For Animal Use Only



Indications: This product has been shown to be effective for the vaccination of healthy cattle five months of age or older against Infectious Bovine Rhinotracheitis and Parainfluenza, Duration of immunity for this product has not been determined. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Directions: Aseptically rehydrate the freeze-dried modified live vaccine with the accompanying bottle of sterile diluent. Mix reconstituted vial well. Administer 2.0 mL subcutaneously in the side of the neck.

Cautions: Do not vaccinate pregnant cows. Do not use in calves nursing pregnant cows. Abortions may result. Store at 2° to 8°C. Use entire contents when first opened. Inactivate unused contents before disposal. Do not vaccinate within 21 days of slaughter. Do not mix with other products, except as specified on this label. In case of human exposure, contact a physician. This product contains Neomycin and Nystatin as preservatives. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitized, are stressed due to shipment or environmental

conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.